

*G. & H. Hobbs to Weymouth
Georgia May 1941*

RING-MASTER

PATENT PENDING



The Superior

Composition Book

Type - locality 8

11 miles north of Lyons -

P. ~~truncatus~~
C. ~~detritus~~

591.9 Alachua

823.5 waycross - 6 7/10 gals. - #1.44

802.6 Lyons

5-1841-1

809.5 - Leomb - Emmanuel Co. line

813.6 - Pitcher plants field, pines, Erica ulmifolia,
mucky sand. Burrows extremely complex,
one passage to water in which nearly
every instance divided 1 or more times,
below the water table. Amphipods & copepods
were present in burrows & a fly larva
(preserved). More than a dozen burrows
were dug, & each contained numerous
small carfish, approx. 3/4 inch long.

Not a single male found, all burrows
dug within a radius of 4 ft. Water
table about 1 1/2 ft. below surface

Technique for collecting: Burrows were dug

til the water was exposed in the
bottle, then thoroughly slushed,
left alone & revisited in about 5 mins.

Grayfish were found sitting on
water level, with part of body
exposed. Method yielded about
60% average.

Invertebrates $6 \frac{30}{10}$ gals. $\$1.33$

all drinks .30

31
24

5-2541-1

5-2541-2
148.11-

How wide, narrow, the diameter of
cup and gray area is wider than
usual. Disposition of furrow
only. Burrows with numerous
a-f-p-h. I is up about 3 inches
above the surface, dip in
Rostelakid of wood in $1 \frac{1}{2}$ inch

Oxistida taken from *Spizella socialis*
↳ Utricularia in shallow pools
at end of culvert. Eox, Vitellina,
Lactaria, P. im., Fundulus shufeldti,
& Gambusia seen in water.

227.5 - High Springs
148.4

79.1